



Instruction Sheet TONSILLECTOMY/ ADENOIDECTOMY

Baptist Downtown • Southside • Baptist South

Tonsillectomy and adenoidectomy are surgical procedures that are very commonly performed. The most common reasons for this type of surgery include persistent infections or enlargement that obstructs breathing. Although your doctor may choose to hospitalize you overnight for medical reasons, these procedures are also successfully performed on an outpatient basis. We do recommend that you plan to take off at least one full week post-operatively from work or school.

PRE-OPERATIVE INSTRUCTIONS:

- Unless prescribed by a physician, do not take any aspirin, aspirin products or NSAIDs one week prior to surgery or after surgery. **(Be aware that for certain surgeries, your surgeon may request that you stay off these types of drugs longer than one week before or after surgery. Please follow their instructions for your specific surgery).** Tylenol is okay.

IMPORTANT: If your physician has placed you on a daily dosage of aspirin or other blood thinner such as Coumadin or Plavix, check with your prescribing doctor regarding instructions for preoperative and postoperative aspirin use. If you have been placed on aspirin by a cardiologist, please discuss your specific stop and restart times with both your cardiologist and surgeon.

- Have nothing to eat or drink after midnight the day before surgery.
- If you take daily medications, take them the morning of surgery with a sip of water. If you have questions about this, check with the hospital preop nurse during the presurgical evaluation.
- Please make arrangements for transportation to the hospital on the day of surgery.

POST-OPERATIVE INSTRUCTIONS:

What to expect:

- You will experience a significant sore throat immediately after surgery. Typically the pain will improve slowly over the first 7 to 10 days; occasionally, there will be a sudden, usually brief, relapse of pain towards the end of the first week. Throat discomfort will last to some extent for several weeks. Many patients will complain of ear pain; however, this is "referred pain" from the throat and is not indicative of any ear problems. Generally within one week to ten days after surgery, most patients are feeling significantly better.
- You may also notice white patches in the throat or bad breath. These are entirely normal and are indicative of the healing process.
- Nausea and vomiting are rather common and can occur for up to 24 hours after the surgery. If this persists, please call our office.

- Some patients will cough up small amounts of dried blood, particularly within the first 24 hours. However, if there are any episodes of heavy bleeding, apply an ice pack to the neck, chew on ice cubes, and call our office. If the office is closed the answering service will direct your call to your surgeon or the on-call physician for further assistance. If bleeding is severe, go to Baptist Medical Center's emergency room.
- Reflux of food into the nose can sometimes occur after surgery. If it lasts for more than a few weeks, please call the office or bring this to your surgeon's attention during your follow up visit.

- Patients sometimes will experience low-grade fevers after surgery that may persist for one to two days. Temperatures can reach as high as 102 to 104 degrees, but usually will resolve with Tylenol and fluids. If you have a high fever (greater than 101 degrees) that lasts longer than 24 hours without any improvement, you will need to notify the office.

- At any time during the postop period, please call the office if you have any questions or concerns about excessive bleeding, breathing difficulty, pain, persistent fever, nausea, swelling or other concerns that seem out of the ordinary from what you have discussed with your surgeon or read in this handout.

Activity:

- Bed rest and very light activity is the rule for the first 24 hours postoperatively. You may increase your activity level as necessary, but use common sense.

- Avoid any heavy lifting, bending, straining, or stooping for at least 2 weeks after surgery as this will put additional pressure on the operative site and may increase your chances of postoperative bleeding. If you have questions about certain activities, please ask us.

- Try to keep your head elevated during sleep for at least 3 days after surgery. Sleeping on 2 or more pillows is effective. A recliner chair is also an excellent option.

- Depending on the individual and the type of surgery done, disability from work may vary. Most patients are able to return to work or school within 1 full week after surgery.

- It is okay to shower or bathe postoperatively. Avoid particularly hot or steamy showers for several days after surgery.

Diet:

- It cannot be emphasized enough that adequate hydration is crucial. This means plenty of fluids at all times, especially Gator Ade, decaffeinated soft drinks or tea, cool soups, Popsicles, ice cream, and Jell-O. Soft foods may be taken later as tolerated; however, it is not essential to begin taking solid foods right away. Stay away from citrus fruits, acidic juices and hot/spicy foods for about 2 weeks.



Instruction Sheet
TONSILLECTOMY/ ADENOIDECTOMY

Baptist Downtown • Southside • Baptist South

Medication:

- Pain medication such as Tylenol with Codeine or hydrocodone will be provided for relief of pain and this should be taken as necessary. You may also use over-the-counter sprays or lozenges indicated for relief of sore throat. You may take Tylenol or Extra Strength Tylenol as directed if the pain is no longer bad enough to require the prescription medication; however, do not combine this with the prescription pain reliever, which already contains Tylenol. If you (or your child) are unable to take medication by mouth, Tylenol suppositories can be given and these are available over the counter at most local drug stores.

- Tylenol may also be taken for mild fever. If postoperative fever (>101 degrees) persists for more than 24 hours, notify the office.

- Antibiotics may be given during the postoperative period. Take all medications as prescribed for you by the physician. If you are nauseous, you can begin the antibiotics the day after surgery. Call the office for any adverse reactions to your medications (vomiting, diarrhea, rash, difficulty breathing or swallowing).

- Unless aspirin has been prescribed as a daily medication, do not take aspirin or ibuprofen-like products (NSAIDS) as they may cause bleeding. These products include: Ecotrin, Bayer, Bufferin, Excedrin, Alka Seltzer, Goody's powders, Motrin, Nuprin, Advil, Alleve, and Naprosyn.

Wound Care:

- No special care is required for the throat after surgery. If you wish, you may rinse or gargle lightly after several days with baking soda in water.

Follow-up:

- Typically, your surgeon will ask you to call the office to give us a “progress report” several days after surgery. We also request that you call us again about 2 weeks after surgery for an update. Most patients do not need to return to the office to be reexamined after surgery. However, depending on the problems you encounter postop, your physician may request that you do so if he feels that it is prudent.

If you have any questions regarding this procedure please discuss them with your surgeon prior to your operation. It is very important that you fully understand the alternatives, the risks, and the complications prior to signing the consent form. Your signature on the consent form verifies that you have read the above instructions, your questions have been answered, and you wish to proceed with the surgical procedure.